for the referral!

Checklist for Inflectra (infliximab-dyyb) Referral

Required documentation for all initial referrals

Dationt	DOB Date						
Piease i	return completed checklist and checklist items for an infusion referral:						
	Patient demographics (e.g. address, phone number, SSN, etc.)						
	Insurance information and copy of insurance card(s). Please indicate the insurance that is primary, and the insurance that is secondary, if applicable, and the subscriber's date of birth. o If insurance requires prior authorization, please provide the phone number and allow up to 15-30 days for this to be completed by one of our Infusion Coordinators.						
	Signed and completed Inflectra Standard Order (our order form) with ICD diagnosis code O Standard Order forms are available at lowcountryrheumatology.com/infusions/						
	Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy, and how long patient has been on Inflectra.						
	Lab results and/or tests to support diagnosis. Pre-Screening: Required TB screening results: PPD (within 1 year) or QuantiFERON Gold Test (within 3 years) Required Hepatitis screening (within 1 year): Hepatitis B Surface Antigen, Hepatitis B Surface Antibody, or Hepatitis B Core Antibody results Lab results within last 60 days: ESR/CRP (if available) Most Recent Rapid 3 (if available)						
	Please indicate name and direct phone number of a contact within your office that we can speak with to obtain any additional information: O Name:						
	o Phone Number:						
	Paperwork can be faxed to (843)-793-6181						
	Infusion Coordinators can assist you with any questions at (843)-572-8932						
	Low Country Rheumatology Infusion Locations Please mark preferred location and we will do our best to accommodate, however we cannot make any guarantees.						
Summerville 2001 2nd Ave, Suite 201, Summerville, SC 29486							
	Mount Pleasant						
	1100 Johnnie Dodds Blvd, Mt. Pleasant, SC 29464						
	West Ashley 2291 Henry Tecklenburg Drive, Charleston, SC 29414						
docume	untry Rheumatology Infusion Services will complete insurance verification and submit all required clinical entation to the patient's insurance company for eligibility. Our Infusion Coordinators will notify you if any furthe tion is required. The patient will have an annual 30-minute consult with our NP to obtain H&P for chart. We wi						

Low Country Rheumatology Use Only Existing Patient Yes____ No____ Physician _____

review financial responsibility with the patient and refer them to any available co-pay assistance as required. Thank you

Standard Orders for Inflectra (infliximab-dyyb) Administration

Patient		DOB	Date	!	_	
*NOTE: Patient is ineligible to re		•	-	_		
process due to the possibility of	f developing a super infe	ction related t	o its effect on the im	mune syste	m.	
Indication: Please indicate the I	highest level of specificity	<i>'</i> .				
□ K50.0 Crohn's		□ K51.9	_Ulcerative Colitis,	□ K60.3	Anal Fistula	
Disease (small intestine)		Unspecified				
□ K50.1Crohn's	□ K51.5	_Left-sided	□ Other ICD-10 Code			
Disease (large intestine)	_	hronic) Colitis				
□ K50.8 Crohn's			_ Other Ulcerative			
Disease (small and large		(chronic) Col	ITIS			
intestine) □ K63.2 Fistula of		□ K51.0	Universal			
intestine			hronic) Pancolitis			
History:		Oleciative (e	in onic, i unconcis	_		
□ Inadequate response to DMA	.RD		□ Unable to tolerat	e DMARD		
□ Rapid 3		□ Swollen/tender joints				
□ ESR/CRP						
·	CAL		□ Progressive erosive arthropathy			
□HBsAg, HBsAb, HB core Ab, H0	LAD	□ Recent of upcom	Recent or upcoming surgery			
☐ History of skin cancer						
 Obtain patient weight Evaluate patient for acconcerns as noted on I Baseline vitals will be obe obtained more freq Titrate infusion over 2 to previous infusion re If infusion reaction occand Procedure Manua Discharge instructions Dose: Inflectra (infliximab- Frequency: Initiation of Inflectra Maintenance dose e 	p, HB core Ab, HCAb negal each visit stive infections, prior or ultifusion Record obtained prior to adminisquently if patient's conditional hours as recommended it action. After dose 4, titracurs, slow or stop infusional. to include possible infusional dyyb)mg/kg in to be administered at well as to be administered at well as the single prior of the single p	stration, hourlion warrants in Pfizer Infusiate infusion over and initiation side effect	y during infusion and t. on guide for doses 1 er 1 hour as tolerate e infusion reaction p s and follow-up appo	at the end -4, and for p d. rotocol per	rgies, or any current health of the infusion. Vital signs will patients receiving pre-med due Articularis Healthcare Policy medule	
Premedicate:						
□ No pre-med						
□ Pre-medicate x 1 dose 30 mir	· ·		405 0 1 44 1	L 11 / 0 ·		
□ 1000 mg Acetamino	pnen PO 🗆 25mg Ben	nadryi PO/IV	□ 125mg Solu-Medi	rollv 🗆 Ot	:her	
Additional orders/comment						
Dractice Name:						
Practice Name:		NPI:				
Physician Name:			State License:			
Physician Signature:						
Date:		UPIN:				