for the referral!

Checklist for Renflexis (infliximab-abda) Referral

Required documentation for all initial referrals

Dationt	DOB Date Dew Start Maintenance					
Please i	return completed checklist and checklist items for an infusion referral:					
	Patient demographics (e.g. address, phone number, SSN, etc.)					
	Insurance information and copy of insurance card(s). Please indicate the insurance that is primary, and the insurance that is secondary, if applicable, and the subscriber's date of birth. o If insurance requires prior authorization, please provide the phone number and allow up to 15-30 days for this to be completed by one of our Infusion Coordinators.					
	Signed and completed Renflexis Standard Order (our order form) with ICD diagnosis code o Standard Order forms are available at lowcountryrheumatology.com/infusions/					
	Supporting clinical MD notes to include any past tried and/or failed therapies, intolerance, outcomes or contraindications to conventional therapy, and how long patient has been on Renflexis.					
	Lab results and/or tests to support diagnosis. Pre-Screening: Required TB screening results: PPD (within 1 year) or QuantiFERON Gold Test (within 3 years) Required Hepatitis screening (within 1 year): Hepatitis B Surface Antigen, Hepatitis B Surface Antibody, or Hepatitis B Core Antibody results Lab results within last 60 days: ESR/CRP (if available) Most Recent Rapid 3 (if available)					
	Please indicate name and direct phone number of a contact within your office that we can speak with to obtain any additional information: O Name:					
	o Phone Number:					
	Paperwork can be faxed to (843)-793-6181					
	Infusion Coordinators can assist you with any questions at (843)-572-8932					
	Low Country Rheumatology Infusion Locations Please mark preferred location and we will do our best to accommodate, however we cannot make any guarantees.					
	Summerville 2001 2nd Ave, Suite 201, Summerville, SC 29486					
	Mount Pleasant					
	1100 Johnnie Dodds Blvd, Mt. Pleasant, SC 29464					
	West Ashley 2291 Henry Tecklenburg Drive, Charleston, SC 29414					
docume	untry Rheumatology Infusion Services will complete insurance verification and submit all required clinical entation to the patient's insurance company for eligibility. Our Infusion Coordinators will notify you if any further tion is required. The patient will have an annual 30-minute consult with our NP to obtain H&P for chart. We will					

Low Country Rheumatology Use Only Existing Patient Yes____ No____ Physician _____

review financial responsibility with the patient and refer them to any available co-pay assistance as required. Thank you

Standard Orders for Renflexis (infliximab-abda) Administration

Patient	DOB	Da	ate								
*NOTE: Patient is ineligible to receive Renfle				ing antibiotic for active							
infectious process due to the possibility of d	eveloping a super infecti	on related to its eff	ect on the im	nmune system.							
Indication: Please indicate the highest level	of specificity.										
☐ K50.0 Crohn's Disease (small	□ K51.9Ulcerativ	e Colitis,	□ K60.3	Anal Fistula							
intestine)	Unspecified										
☐ K50.1Crohn's Disease (large				D-10 Code							
intestine)	(chronic) Colitis										
☐ K50.8 Crohn's Disease (small and	□ K51.8 Other Ul	cerative (chronic)									
large intestine)	Colitis □ K51.0 Universal Ulcerative										
☐ K63.2 Fistula of intestine	☐ K51.0Universa (chronic) Pancolitis	l Olcerative									
	(cirrorne) Fancontis										
History:											
□ Inadequate response to DMARD		□ Unable to tole									
□ Rapid 3		☐ Swollen/tender joints									
□ ESR/CRP		☐ Progressive erosive arthropathy									
□HBsAg, HBsAb, HB core Ab, HCAb		☐ Recent or upco	Recent or upcoming surgery								
☐ History of skin cancer											
Orders:											
□ Standard Order Protocol:											
 Confirm current PPD, Tspot, or CXR 	1										
Confirm HbsAg, HBsAb, HB core Ab, HCAb negative											
 Obtain patient weight each visit 											
 Evaluate patient for active infection 	ns, prior or upcoming sur	gical procedures, m	nedication all	ergies, or any current health							
concerns as noted on Infusion Reco	· · · · · · · · · · · · · · · · · · ·	,		,							
 Baseline vitals will be obtained prior to administration, hourly during infusion and at the end of the infusion. Vital signs will be obtained more frequently if patient's condition warrants it. Titrate infusion over 2 hours as recommended in Pfizer Infusion guide for doses 1-4, and for patients receiving pre-med du to previous infusion reaction. After dose 4, titrate infusion over 1 hour as tolerated. If infusion reaction occurs, slow or stop infusion, and initiate infusion reaction protocol per Articularis Healthcare Policy 											
							and Procedure Manual.	stop iniusion, and initia	te illusion reaction	ii protocoi pe	a Articularis ricularicare rolley
							Discharge instructions to include por	ossible infusion side effe	rts and follow-up a	nnointment s	chedule
								ossible illiasion side effec	co and renew up a	oponitinent s	circulate
							Dose:				
□ Renflexis (infliximab-abda)	mg/kg in Normal Sali	ne IV									
Frequency:											
☐ Initiation of Renflexis to be admir	nistered at week(s) 0, 2, a	and 6									
□ Maintenance dose every	_ weeks										
Premedicate:											
□ No pre-med											
□ Pre-medicate x 1 dose 30 minutes prior to	each infusion with:										
•	□ 25mg Benadryl PO/IV	□ 125mg Solu-M	edrol IV 🗆 (Other							
	. 6 , ,	. 0									
Additional orders/comments:											
Practice Name:		NDI.									
		NPI:									
Physician Name:		State License:									
Physician Signature:		DEA #:									
Date:											
246.		I IDINI:									